arXiv:1608.01051v1 [math.OA] 3 Aug 2016

ROKHLIN DIMENSION OF $\mathbb{Z}^m$-ACTIONS ON SIMPLE C^*-ALGEBRAS

HUNG-CHANG LIAO

Abstract. We study Rokhlin dimension of $\mathbb{Z}^m$-actions on simple separable stably finite nuclear C^*-algebras. We prove that under suitable assumptions, a strongly outer $\mathbb{Z}^m$-action has finite Rokhlin dimension. This extends the known result for automorphisms. As an application, we show that for a large class of C^*-algebras, the $\mathbb{Z}^m$-Bernoulli action has finite Rokhlin dimension.

1. Introduction

Extending topological covering dimension to the context of nuclear C^*-algebras, Winter and Zacharias introduced *nuclear dimension* in [35]. Recent advances in the study of nuclear C^*-algebras have revealed that:

(1) Finiteness of nuclear dimension implies other structural properties, including Jiang-Su stability and strict comparison ([34, 28]). In fact, a conjecture of Toms and Winter predicts that these are all equivalent for unital separable simple nuclear C^*-algebras.
(2) Unital separable simple C^*-algebras of finite nuclear dimension have been completely classified by K-theoretic invariants, provided they satisfy a technical condition known as the UCT (Universal Coefficient Theorem) ([9, 26, 32, 4]).

Given the success above, it is important to understand how nuclear dimension behaves under various constructions, such as crossed products. Motivated by the Kakutani-Rokhlin lemma in ergodic theory ([8, 27]), its non-commutative analogues for finite von Neumann algebras ([2, 22]) and later developments in the C^*-realm, Hirshberg, Winter, and Zacharias introduced *Rokhlin dimension* in [6] for actions of finite groups and the set of integers $\mathbb{Z}$ on unital C^*-algebras. One of the main results in [6] is that finiteness of nuclear dimension passes to crossed products when the action has finite Rokhlin dimension. In addition, they showed that:

(1) Actions with finite Rokhlin dimension are ubiquitous; more precisely, given a unital Jiang-Su stable C^*-algebra A, the automorphisms with finite Rokhlin dimension form a G_δ-subset of the automorphism group $\mathrm{Aut}(A)$ ([6, Theorem 3.4]).
(2) There are plenty of natural examples coming from actions on commutative C^*-algebras, i.e., topological dynamics. In fact, given any minimal homeomorphism of a compact metrizable space with finite covering dimension, the induced C^*-system has finite Rokhlin dimension ([6, Theorem 6.1]).

The notion of Rokhlin dimension has been generalized to actions of $\mathbb{Z}^m$ by Szabó [30], and more generally to actions of residually finite groups by Szabó, Wu, and Zacharias [31]. Similar results regarding nuclear dimension, genericity, and topological dynamics were also obtained.

The present paper focuses on C^*-dynamical systems of a purely non-commutative nature. We consider $\mathbb{Z}^m$-actions on simple separable stably finite nuclear (hence tracial) C^*-algebras. Our first main result is a Rokhlin type theorem: under suitable assumptions, a certain strong form of outerness, which is a necessary condition for the action to have finite Rokhlin dimension (see Proposition 2.3), turns out to be sufficient. The precise statement is given below (see Section 2 and Section 5 for relevant definitions) Recall that *Property (SI)*, as introduced by Sato in [29], is a comparison-by-traces type property for central sequence algebras (see Definition 5.2 for the precise definition). This property is possessed by a large class of simple nuclear C^*-algebras, in particular the ones of finite nuclear dimension (cf. [18, 34, 28]).

Theorem 1.1. *(Theorem 5.5) Let A be a unital simple separable nuclear C^*-algebra with nonempty trace space $T(A)$, and let $\alpha : \mathbb{Z}^m \to \mathrm{Aut}(A)$ be a group action. Suppose*

(1) *A has property (SI),*
(2) *α is strongly outer,*
(3) *$T(A)$ is a Bauer simplex with finite dimensional extreme boundary,*
(4) *$\tau \circ \alpha = \tau$ for every $\tau \in T(A)$.*

Then $\dim_{\mathrm{Rok}}(\alpha) \leq 4^m - 1$.

This type of results dates back to Connes' non-commutative Rokhlin lemma [3], where he showed that outer $\mathbb{Z}$-actions on finite von Neumann algebras have the (W^*)-Rokhlin property. Ocneanu in [22] obtained a similar result for actions of discrete amenable groups. On the C^*-side, Hermann and Ocneanu introduced a Rokhlin property for automorphisms of C^*-algebras [5]. The definition was later refined by Kishimoto [13], who proved that for certain AF and A$\mathbb{T}$ algebras, strong outerness implies the Rokhlin property [14, 15]. Nakamura studied $\mathbb{Z}^2$-actions on UHF algebras and obtained a similar Rokhlin type result [20]. Various Rokhlin type definitions and theorems also appeared in the work of Osaka and Phillips [24, 23], Sato [29], and Matui and Sato [17, 19]. Recently, we proved a Rokhlin type theorem similar to Theorem 1.1 for automorphisms [16], which was built upon a remarkable breakthrough made by Matui and Sato (unpublished; see [16, Section 6]).

In Section 2 we will prove the converse of Theorem 1.1 (see Proposition 2.3). Combining this and Remark 2.4, we obtain the following corollary:

Corollary 1.2. *Let A be a unital simple separable nuclear C^*-algebra with nonempty trace space $T(A)$, and let $\alpha : \mathbb{Z}^m \to \mathrm{Aut}(A)$ be a group action. Suppose*

(1) *A has property (SI),*
(2) *$T(A)$ is a Bauer simplex with finite dimensional extreme boundary, and*
(3) *$\tau \circ \alpha = \tau$ for every $\tau \in T(A)$.*

Then the following are equivalent.

(i) $\dim_{Rok}(\alpha) \leq 4^m - 1$.
(ii) *For every $v \in \mathbb{Z}^m$, the restriction map $r : T(A \rtimes_{\alpha^v} \mathbb{Z}^m) \to T(A)^{\alpha^v}$ is injective.*
(iii) *The action α is strongly outer.*

While in general it is not easy to verify that a given action has finite Rokhlin dimension, strong outerness can often be checked explicitly. As an example, we study Bernoulli shifts in Section 6 and make the following observation.

Proposition 1.3. *(Proposition 6.1) Let A be a unital simple C^*-algebra with a unique trace. Suppose A is not isomorphic to the algebra of complex numbers $\mathbb{C}$. Then the Bernoulli action $\sigma : \mathbb{Z}^m \to \mathrm{Aut}(\bigotimes_{\mathbb{Z}^m} A)$ is strongly outer. In particular, σ has finite Rokhlin dimension.*

In addition to this proposition, we observe that the Rokhlin dimension of a Bernoulli shift can be bounded by the Rokhlin dimension of a subshift (Proposition 6.3). Combining these two results, we obtain a large class of examples of $\mathbb{Z}^m$-actions with finite Rokhlin dimension.

Let us describe how the paper is organized. In Section 2 we review the definition of Rokhlin dimension for $\mathbb{Z}^m$-actions. We show that as in the case of automorphisms, strong outerness is a necessary condition for an $\mathbb{Z}^m$-action to have finite Rokhlin dimension. We also review the notion of continuous W^*-bundles and discuss group actions on them. In Section 3 we prove a Rokhlin type theorem for $\mathbb{Z}^m$-actions on continuous W^*-bundles. The ideas and techniques are almost identical to the case of automorphisms, hence we will only outline the important steps and we refer the reader to [16] for full details. Section 4 is devoted to the technical notion we call the "approximate Rokhlin property", which already appeared in our previous paper. Here it serves the same purpose: a bridge connecting the Rokhlin property at the level of W^*-bundles and Rokhlin dimension at the level of C^*-algebras. Section 5 contains a simple but crucial observation (Proposition 5.1), which allows us to obtain finite Rokhlin dimension for $\mathbb{Z}^m$-actions from "equivariant Rokhlin towers for automorphisms" This section also contains a technical result which establishes the existence of these Rokhlin towers. Combining the observations and results from previous sections, we establish Theorem 1.1. In the final section, Section 6, we study Bernoulli shifts on infinite tensor products. Combining the Rokhlin type theorem from Section 5 and an observation involving subshifts, we show that for a large class of C^*-algebras, the $\mathbb{Z}^m$-Bernoulli shift has finite Rokhlin dimension.

Acknowledgment: Much of this work was done under the Pritchard Dissertation Fellowship granted by the Department of Mathematics at Pennsylvania State University. The author wishes to thank the Department of Mathematics for its support. The author also wishes to thank his dissertation advisor, Nate Brown, for many valuable discussions, including the suggestion of Proposition 6.3.

2. Preliminaries

Let us first fix some notations. Let A be a C^*-algebra, Γ be a discrete group, and $\alpha : \Gamma \to \mathrm{Aut}(A)$ be a group action. We write α^g for the automorphism $\alpha(g) \in \mathrm{Aut}(A)$. Suppose

the trace space $T(A)$ of A is nonempty, then the set of α^g-invariant traces will be denoted $T(A)^{\alpha^g}$. Given any $\tau \in T(A)$, we write $(\pi_\tau, \mathcal{H}_\tau)$ for the GNS representation associated to τ.

Definition 2.1. ([17, Definition 2.7]) Let A be a unital C^*-algebra with nonempty trace space $T(A)$, and let Γ be a discrete group. An action $\alpha : \Gamma \to \mathrm{Aut}(A)$ is called *strongly outer* if for every $g \in \Gamma \setminus \{e\}$ and every $\tau \in T(A)^{\alpha^g}$, the weak extension $\tilde{\alpha}^g \in \mathrm{Aut}(\pi_\tau(A)'')$ is an outer automorphism.

From now on we focus on the case $\Gamma = \mathbb{Z}^m$. We use $\xi_1, ..., \xi_m$ for the standard generators of $\mathbb{Z}^m$ (i.e., ξ_i is the unit vector with 1 on the i-th place and 0's elsewhere). The automorphisms α^{ξ_i} will be denoted α^i for short. For each $n \in \mathbb{N}$, we write B_n^m for the "box" $\{0, 1, ..., n-1\}^m$ in $\mathbb{Z}^m$. When there is no possible confusion, we drop the dimension m and simply write B_n.

Below we recall the definition of Rokhlin dimension for $\mathbb{Z}^m$-actions. Roughly speaking, an action has finite Rokhlin dimension if we can find families of positive contractions in the algebra which are almost central, almost cyclically permuted by the action, almost mutually orthogonal, and almost sum to 1.

Definition 2.2. ([30, Definition 1.6]) Let A be a unital C^*-algebra, and let $\alpha : \mathbb{Z}^m \to \mathrm{Aut}(A)$ be a group action. We say that the action α has Rokhlin dimension at most d, and write $\dim_{\mathrm{Rok}}(\alpha) \leq d$, if for any finite subset $F \subseteq A$, $\varepsilon > 0$, and $n \in \mathbb{N}$, there exist positive contractions

$$(f_v^{(\ell)})_{v \in B_n}^{\ell = 0, 1, ..., d}$$

in A satisfying the following properties:

(1) $\|\alpha^w(f_v^{(\ell)}) - f_{v+w}^{(\ell)}\| < \varepsilon \mod n \qquad (0 \leq \ell \leq d,\ v \in B_n,\ w \in \mathbb{Z}^m)$.
(2) $\|f_v^{(\ell)} f_{v'}^{(\ell)}\| < \varepsilon \qquad (0 \leq \ell \leq d,\ v, v' \in B_n,\ v \neq v')$.
(3) $\|[f_v^{(\ell)}, a]\| < \varepsilon \qquad (0 \leq \ell \leq d,\ v \in B_n,\ a \in F)$.
(4) $\left\| \sum_{\ell=0}^d \sum_{v \in B_n} f_v^{(\ell)} - 1_A \right\| < \varepsilon$.

As in the case of $\mathbb{Z}$-actions, strong outerness is a necessary condition for an action to have finite Rokhlin dimension (when the C^*-algebra has traces).

Proposition 2.3. *Let A be a unital C^*-algebra with nonempty trace space $T(A)$, and let $\alpha : \mathbb{Z}^m \to \mathrm{Aut}(A)$ be a group action. If $\dim_{\mathrm{Rok}}(\alpha) = d < \infty$, then α is strongly outer.*

Proof. Since the argument is almost identical to the one for implication (1) $\implies$ (3) of [16, Corollary 1.2], we only sketch the proof here. Given $v \in \mathbb{Z}^m$, let u be the implementing unitary in the crossed product $A \rtimes_{\alpha^v} \mathbb{Z}$. Given any tracial state φ on $A \rtimes_{\alpha^v} \mathbb{Z}$, one can show that $\varphi(au^n) = 0$ for any $a \in A$ and $n \in \mathbb{N}$ by approximating 1_A by the sum of the Rokhlin elements. Then the proof of [14, Lemma 4.4] shows that a weak extension of α^v cannot be inner. □

Remark 2.4. Note that the argument above also shows that for each $v \in \mathbb{Z}^m$, the restriction map $r : T(A \rtimes_{\alpha^v} \mathbb{Z}^m) \to T(A)^{\alpha^v}$ is injective (hence bijective).

We briefly review some basic definitions of continuous W^*-bundles, which was introduced by Ozawa in [25] (see also [1]). A *continuous W^*-bundle* consists of the following data:

- a C^*-algebra $\mathcal{M}$,
- a compact metrizable space K such that $C(K)$ sits inside the center of $\mathcal{M}$, and
- a faithful conditional expectation $E : \mathcal{M} \to C(K)$ satisfying the tracial condition $E(ab) = E(ba)$ for all $a, b \in \mathcal{M}$,

such that the norm-closed unit ball of $\mathcal{M}$ is complete in the *uniform 2-norm* $\|\cdot\|_{2,u}$, defined by

$$\|x\|_{2,u} := \|E(x^*x)\|^{1/2}.$$

For each $\lambda \in K$, the GNS representation $\pi_\lambda(\mathcal{M})$ coming from the tracial state $\tau_\lambda := ev_\lambda \circ E$ is called a *fiber* of the bundle $(\mathcal{M}, K, E)$.

Given a unital simple separable C^*-algebra with nonempty trace space $T(A)$, we define the *uniform 2-norm* $\|\cdot\|_{2,u}$ on A by

$$\|a\|_{2,u} := \sup_{\tau \in T(A)} \tau(a^*a)^{1/2}.$$

Let $\overline{A}^u$ be the completion of A with respect to $\|\cdot\|_{2,u}$. Ozawa proved in [25] that when $T(A)$ is a *Bauer simplex*, i.e., when the extremem boundary $\partial_e T(A)$ is compact, the set of continuous functions on $\partial_e T(A)$ can be embedded into the center of $\overline{A}^u$. Moreover, there is a faithful tracial conditional expectation

$$E : \overline{A}^u \to C(\partial_e T(A))$$

satisfying $E(a)(\tau) = \tau(a)$ for each $a \in A$ and $\tau \in \partial_e T(A)$. Therefore $(\overline{A}^u, \partial_e T(A), E)$ has the structure of a continuous W^*-bundle. In this case each fiber $\pi_\tau(\overline{A}^u)$ is isomorphic to the weak closure $\pi_\tau(A)''$.

Given a free ultrafilter ω on $\mathbb{N}$, one can form the *ultrapower* $(\mathcal{M}^\omega, E^\omega, K^\omega)$ and the *central sequence W^*-bundle* $(\mathcal{M}^\omega \cap \mathcal{M}', E^\omega, K^\omega)$ of a continuous W^*-bundle $(\mathcal{M}, K, E)$ in a way similar to the case of tracial von Neumann algebras. The uniform 2-norm on $\mathcal{M}^\omega$ will be denoted $\|\cdot\|_{2,u}^{(\omega)}$. For details, we refer the reader to [1] and [16].

Now we discuss group actions on continuous W^*-bundles. Let $(\mathcal{M}, K, E)$ be a continuous W^*-bundle and let Γ be a discrete group. A Γ-*action* on $(\mathcal{M}, K, E)$ consists of a pair (β, σ), where

$$\beta : \Gamma \to \mathrm{Aut}(\mathcal{M}) \qquad \text{and} \qquad \sigma : \Gamma \to \mathrm{Homeo}(K)$$

are group homomorphisms, such that the diagram

$$\begin{array}{ccc} \mathcal{M} & \xrightarrow{\beta^g} & \mathcal{M} \\ {\scriptstyle E}\downarrow & & \downarrow{\scriptstyle E} \\ C(K) & \xrightarrow{\bar{\sigma}^g} & C(K) \end{array}.$$

commutes for every $g \in \Gamma$. Given a group action $(\beta, \sigma) : \Gamma \to \mathrm{Aut}(\mathcal{M}, K, E)$, for each $g \in \Gamma$ and $\lambda \in K$, there is a well-defined *-isomorphism

$$\begin{aligned} \beta^g_\lambda : \pi_\lambda(\mathcal{M}) &\to \pi_{(\sigma_g)^{-1}(\lambda)}(\mathcal{M}), \\ \pi_\lambda(x) &\mapsto \pi_{(\sigma_g)^{-1}(\lambda)}(\beta^g(x)). \end{aligned}$$

When σ is the trivial action on K, this gives rise to a group action

$$\beta_\lambda : \Gamma \to \mathrm{Aut}(\pi_\lambda(\mathcal{M})).$$

Let ω be a free ultrafilter on $\mathbb{N}$. A group action (β, σ) can be promoted to an action on the ultrapower or the central sequence W^*-bundle. More precisely, for each $g \in \Gamma$ we have automorphisms

$$\begin{aligned} \beta^g_\omega : \mathcal{M}^\omega &\to \mathcal{M}^\omega, \\ (x_n)_n &\mapsto (\beta^g(x_n))_n, \end{aligned}$$

and

$$\begin{aligned} \bar{\sigma}^g_\omega : C(K^\omega) &\to C(K^\omega), \\ (f_n)_n &\mapsto (f_n \circ \sigma_g)_n. \end{aligned}$$

Therefore we obtain actions

$$\beta_\omega : \Gamma \to \mathrm{Aut}(\mathcal{M}^\omega) \qquad \text{and} \qquad \bar{\sigma}_\omega : \Gamma \to \mathrm{Aut}(C(K^\omega)).$$

One easily checks that the diagram

$$\begin{array}{ccc} \mathcal{M}^\omega & \xrightarrow{\beta^g_\omega} & \mathcal{M}^\omega \\ {\scriptstyle E^\omega}\downarrow & & \downarrow{\scriptstyle E^\omega} \\ C(K^\omega) & \xrightarrow{\bar{\sigma}^g_\omega} & C(K^\omega) \end{array}$$

commutes for every $g \in \Gamma$. We write $(\mathcal{M}^\omega \cap \mathcal{M}')^{\beta_\omega}$ for the fixed point subalgebra of the central sequence W^*-bundle $\mathcal{M}^\omega \cap \mathcal{M}'$.

3. W^*-Rokhlin property

The goal of this section is to prove a Rokhlin type theorem for $\mathbb{Z}^m$-actions on continuous W^*-bundles. Since the argument is almost the same as [16, Theorem 4.1], we will be terse for most of the time. Our first objective is to embed arbitrarily large matrices into the fixed point algebra of the central sequence W^*-bundle, as in the following proposition and corollary. Recall that a completely positive (c.p.) map $\varphi : A \to B$ between C^*-algebras is called *order zero* if $\varphi(a)\varphi(b) = 0$ for all positive elements $a, b \in A$ satisfying $ab = 0$.

Proposition 3.1. *(cf.* [16, Proposition 3.4]*) Let $(\mathcal{M}, K, E)$ be a separable continuous W^*-bundle and $(\beta, \sigma) : \mathbb{Z}^m \to \mathrm{Aut}(\mathcal{M}, K, E)$ be a group action. Suppose*

(1) *the action is trivial on K, i.e., $\sigma = \mathrm{id}_K$;*
(2) $\dim(K) = d < \infty$*;*
(3) *each fiber $\pi_\lambda(\mathcal{M})$ is isomorphic to the hyperfinite II_1 factor $\mathcal{R}$;*
(4) *the fiber action $\beta : \mathbb{Z}^m \to \mathrm{Aut}(\pi_\lambda(\mathcal{M}))$ is pointwise outer for each $\lambda \in \mathbb{K}$.*

Then for each $p \in \mathbb{N}$, there exist completely positive contractive (c.p.c.) order zero maps $\psi_1, ..., \psi_{d+1} : M_p(\mathbb{C}) \to (\mathcal{M}^\omega \cap \mathcal{M}')^{\beta_\omega}$ with commuting ranges such that $\psi_1(1) + \cdots + \psi_{m+1}(1) = 1_{\mathcal{M}}$.

Corollary 3.2. *(cf.* [16, Proposition 3.3]*) Let $(\mathcal{M}, K, E)$ be a separable continuous W^*-bundle and $(\beta, \sigma) : \mathbb{Z}^m \to \mathrm{Aut}(\mathcal{M}, K, E)$ be a group action. Suppose*

(1) *the action is trivial on K, i.e., $\sigma = \mathrm{id}_K$;*
(2) $\dim(K) = d < \infty$*;*
(3) *each fiber $\pi_\lambda(\mathcal{M})$ is isomorphic to the hyperfinite II_1 factor $\mathcal{R}$;*
(4) *the fiber action $\beta : \mathbb{Z}^m \to \mathrm{Aut}(\pi_\lambda(\mathcal{M}))$ is pointwise outer for each $\lambda \in \mathbb{K}$.*

*Then for every $p \in \mathbb{N}$, there is a unital *-homomorphism $\sigma : M_p(\mathbb{C}) \to (\mathcal{M}^\omega \cap \mathcal{M}')^{\beta_\omega}$.*

Proof. This follows immediately from Proposition 3.1 and [10, Lemma 7.6]. □

To prove Proposition 3.1, we make use of the finite dimensionality of K and build a collection of c.p.c. maps from a matrix algebra $M_p(\mathbb{C})$ into $\mathcal{M}$ which are almost central, almost invariant under the action β, and have almost commuting ranges. The following lemma is simply a $\mathbb{Z}^m$-version of [16, Lemma 3.8] (with identical proof).

Lemma 3.3. *Under the same assumptions as Proposition 3.1, for every $\varepsilon > 0, p \in \mathbb{N}$, and norm bounded $\|\cdot\|_{2,u}$-compact subset $\Omega \subseteq \mathcal{M}$, there exist c.p.c. maps $\psi^{(1)}, ..., \psi^{(d+1)} : M_p(\mathbb{C}) \to \mathcal{M}$ (remember that $d = \dim(K)$) such that*

(i) $\|[\psi^{(\ell)}(e), x]\|_{2,u} < \varepsilon \qquad (1 \le \ell \le d+1,\ e \in M_p(\mathbb{C})_1,\ x \in \Omega)$*;*
(ii) $\|[\psi^{(\ell)}(e), \psi^{(k)}(f)]\|_{2,u} < \varepsilon \qquad (1 \le \ell \ne k \le d+1,\ e, f \in M_p(\mathbb{C})_1)$*;*
(iii) $\|\psi^{(\ell)}(1)\psi^{(\ell)}(e^*e) - \psi^{(\ell)}(e)^*\psi^{(\ell)}(e)\|_{2,u} < \varepsilon \qquad (1 \le \ell \le d+1,\ e \in M_p(\mathbb{C})_1)$*;*
(iv) $\sum_{\ell=1}^{d+1} \psi^{(\ell)}(1) = 1_{\mathcal{M}}$*;*
(v) $\|\beta^i(\psi^{(\ell)}(e)) - \psi^{(\ell)}(e)\|_{2,u} < \varepsilon \qquad (1 \le i \le m,\ 1 \le \ell \le d+1,\ e \in M_p(\mathbb{C})_1)$.

Now the proof of Proposition 3.1 can be completed by mapping sequences of c.p.c. maps as found in Lemma 3.3 into the ultrapower:

Proof. (of Proposition 3.1) Let $\{x_1, x_2, ...\}$ be a countable $\|\cdot\|_{2,u}$-dense subset of $\mathcal{M}$. By Lemma 3.3, for each $n \in \mathbb{N}$ there exist c.p.c. maps $\psi_n^{(1)}, \psi_n^{(2)}, ..., \psi_n^{(d+1)} : M_p(\mathbb{C}) \to \mathcal{M}$ such that (ii) to (v) in Lemma 3.3 hold with $\varepsilon = \frac{1}{n}$ and

(i') $\|[\psi_n^{(\ell)}(e), x_j]\|_{2,u} < \frac{1}{n} \qquad (1 \le \ell \le d+1,\ e \in M_p(\mathbb{C})_1,\ 1 \le j \le n)$.

For each $1 \le \ell \le d+1$ define c.p.c. maps $\psi_\ell : M_p(\mathbb{C}) \to \mathcal{M}^\omega$ by

$$\psi_\ell(e) = \pi_\omega(\psi_1^{(\ell)}(e), \psi_2^{(\ell)}(e), \psi_3^{(\ell)}(e), ...),$$

where π_ω is the quotient map from $\ell^\infty(\mathbb{N}, \mathcal{M})$ onto $\mathcal{M}^\omega$. Then by construction

- the image of ψ_ℓ belongs to $(\mathcal{M}^\omega \cap \mathcal{M}')^{\beta_\omega}$;
- $\psi_1, \psi_2, ..., \psi_{d+1}$ have commuting ranges;
- $\psi_1(1) + \cdots + \psi_{d+1}(1) = 1_{\mathcal{M}}$;
- $\psi_\ell(1)\psi_\ell(e^*e) = \psi_\ell(e)^*\psi_\ell(e) \qquad (1 \leq \ell \leq d+1,\ e \in M_p(\mathbb{C}))$.

It remains to show that each ψ_ℓ is order zero, but this follows from the fourth bullet (see [16, Lemma 3.9]). □

Now we define the W^*-Rokhlin property and state the Rokhlin type theorem.

Definition 3.4. Let $(\mathcal{M}, K, E)$ be a separable continuous W^*-bundle and $(\beta, \sigma) : \mathbb{Z}^m \to \mathrm{Aut}(\mathcal{M}, K, E)$ be a group action. We say (β, σ) has the *W^*-Rokhlin property* if for every $n \in \mathbb{N}$ there exist projections $\{p_v\}_{v \in B_n}$ in $\mathcal{M}^\omega \cap \mathcal{M}'$ such that

(1) $(\beta_\omega)^u(p_v) = p_{v+u} \mod n \qquad (v \in B_n,\ u \in \mathbb{Z}^m)$.
(2) $\sum_{v \in B_n} p_v = 1_{\mathcal{M}}$.
(3) $\|1_{\mathcal{M}} - p_0\|_{2,u}^{(\omega)} < 1$.

Theorem 3.5. *Let $(\mathcal{M}, K, E)$ be a separable continuous W^*-bundle and $(\beta, \sigma) : \mathbb{Z}^m \to \mathrm{Aut}(\mathcal{M}, K, E)$ be a group action. Suppose*

(1) *the action is trivial on K, i.e., $\sigma = \mathrm{id}_K$;*
(2) *$\dim(K) = d < \infty$;*
(3) *each fiber $\pi_\lambda(\mathcal{M})$ is isomorphic to the hyperfinite II_1 factor $\mathcal{R}$;*
(4) *the fiber action $\beta : \mathbb{Z}^m \to \mathrm{Aut}(\pi_\lambda(\mathcal{M}))$ is pointwise outer for each $\lambda \in K$.*

Then (β, σ) has the W^*-Rokhlin property.

The idea of proving Theorem 3.5 is the same as [16, Theorem 4.1]. We first obtain a partition of unity which are almost mutually orthogonal in terms of the uniform 2-norm, and then glue the Rokhlin tower in each fiber (the fiberwise result was due to Ocneanu [21]) to a global section. The following lemma provides the partition of unity we need.

Lemma 3.6. *Let $(\mathcal{M}, K, E)$ be a separable continuous W^*-bundle and $(\beta, \sigma) : \mathbb{Z}^m \to \mathrm{Aut}(\mathcal{M}, K, E)$ be a group action. Suppose*

(1) *the action is trivial on K, i.e., $\sigma = \mathrm{id}_K$;*
(2) *for each $p \in \mathbb{N}$ there exists a unital *-homomorphism $\sigma : M_p(\mathbb{C}) \to (\mathcal{M}^\omega \cap \mathcal{M}')^{\beta_\omega}$.*

Then given a norm bounded $\|\cdot\|_{2,u}$-compact subset Ω of $\mathcal{M}$, $\varepsilon > 0$, and a partition of unity $\{g_j\}_{j=1}^N$ subordinate to some open cover of K, there exist positive contractions $a_1, a_2, ..., a_N$ in $\mathcal{M}$ such that

(i) $\|[a_j, x]\|_{2,u} < \varepsilon \qquad (1 \leq j \leq N,\ x \in \Omega)$;
(ii) $\|(\beta_\omega)^i(a_j) - a_j\|_{2,u} < \varepsilon \qquad (1 \leq i \leq m,\ 1 \leq j \leq N)$.
(iii) $\|a_i a_j\|_{2,u} < \varepsilon \qquad (1 \leq i \neq j \leq N)$;
(iv) $\|[a_j^{1/2}, x]\|_{2,u} < \varepsilon \qquad (1 \leq j \leq N,\ x \in \Omega)$;
(v) $\|a_i^{1/2} a_j^{1/2}\|_{2,u} < \varepsilon \qquad (1 \leq i \neq j \leq N)$;
(vi) $|\tau_\lambda(a_j x) - g_j(\lambda)\tau_\lambda(x)| < \varepsilon \qquad (1 \leq j \leq N,\ x \in \Omega,\ \lambda \in K)$;

(vii) $\sum_{j=1}^{N} a_j = 1_{\mathcal{M}}$.

Proof. The proof is identical to [16, Lemma 4.4]. □

Proof. (of Theorem 3.5) The only difference with the proof of [16, Theorem 4.1] is that in order to obtain a Rokhlin tower in each fiber, we invoke Ocneanu's Rokhlin type theorem [21, Theorem 2] instead of Connes' Rokhlin lemma for automorphisms. □

4. Approximate Rokhlin Property

In this section we extend the notion of "invariant σ-ideal" to actions of an arbitrary discrete amenable group. The main result here is Lemma 4.2, which generalizes a similar result for automorphisms and shows that the trace kernel ideal J_A is an invariant σ-ideal under an action of any discrete amenable group.

The notion of σ-ideals was first introduced by Kirchberg and Rørdam in [10]. It was designed to capture the behavior of a quasicentral approximate unit. When B is a C^*-algebra, Γ is a discrete countable amenable group, and $\alpha : \Gamma \to \mathrm{Aut}(B)$ is a group action, we define a notion which we shall call "α-σ-ideal" (cf. [16, Definition 5.5]), which essentially packages an almost invariant quasicentral approximate unit.

Definition 4.1. Let B be a C^*-algebra, Γ a discrete countable amenable group, and $\alpha : \Gamma \to \mathrm{Aut}(B)$ a group action. Let J be a closed ideal in B such that $\alpha^g(J) = J$ for every $g \in \Gamma$. We say J is an *α-σ-ideal* if for every separable C^*-subalgebra C of B satisfying $\alpha^g(C) = C$ for every $g \in \Gamma$, there exists a positive contraction $u \in C' \cap J$ such that

(1) $\alpha^g(u) = u$ for every $g \in \Gamma$.
(2) $uc = c$ for every $c \in C \cap J$.

Let A be a unital separable C^*-algebra and let A_ω be the ultrapower of A (here we fix once and for all a free ultrafilter ω on $\mathbb{N}$). Suppose that A has nonemepty trace space $T(A)$. Given an element a in A_ω represented by a sequence $(a_1, a_2, ...)$, recall from [10] that the seminorm $\|\cdot\|_{2,\omega}$ on A_ω is defined by

$$\|a\|_{2,\omega} := \lim_{n\to\omega} \sup_{\tau\in T(A)} \tau(a_n^* a_n)^{1/2}.$$

The *trace kernel ideal* J_A is defined to be

$$J_A := \{e \in A_\omega : \|e\|_{2,\omega} = 0\}.$$

Lemma 4.2. *Let A be a unital separable C^*-algebra with $T(A) \neq \emptyset$, Γ a discrete countable amenable group, and $\alpha : \Gamma \to \mathrm{Aut}(A)$ a group action. Then the trace-kernel ideal J_A is an α_ω-σ-ideal in A_ω.*

Proof. First note that $(\alpha_\omega)^g(J_A) = J_A$ for every $g \in \Gamma$ since $(\alpha_\omega)^g$ preserves the uniform 2-norm. By [10, Proposition 4.6] J_A is a σ-ideal in A_ω, so there exists a positive contraction $e \in C' \cap J_A$ such that $ec = c$ for every $c \in C \cap J_A$. Let $F_1 \subseteq F_2 \subseteq \cdots \subseteq \Gamma$ be an increasing

sequence of finite subsets whose union is Γ. By amenability for each $n \in \mathbb{N}$ we can find a $(F_n, 1/n)$-invariant (finite) subset K_n of Γ, in the sense that

$$\max_{s\in F_n} \frac{|sK_n \triangle K_n|}{|K_n|} < \frac{1}{n}.$$

For each $n \in \mathbb{N}$, define

$$e^{(n)} := \frac{1}{|K_n|} \sum_{g\in K_n} (\alpha_\omega)^g(e).$$

One checks that $[e^{(n)}, c] = 0$ for every $c \in C$ and $e^{(n)}c = c$ for every $c \in C \cap J$. Moreover, for each $s \in F_n$ we have

$$\begin{aligned} \|(\alpha_\omega)^s(e^{(n)}) - e^{(n)}\| &= \frac{1}{|K_n|} \left\| \sum_{g\in K_n} (\alpha_\omega)^{sg}(e) - \sum_{g\in K_n} (\alpha_\omega)^g(e) \right\| \\ &\leq \frac{1}{|K_n|} |sK_n \triangle K_n| < \frac{1}{n}. \end{aligned}$$

To finish the proof, we invoke Kirchberg's ε-test [10, Lemma 3.1]. Let $d = (d_1, d_2, ...)$ be a strictly positive contraction in the separable C^*-algebra $C \cap J_A$, and $\{g_k\}_{k=1}^\infty$ be a list of elements in Γ. Take a dense sequence $\{c^{(k)}\}_{k=1}^\infty$ in the unit ball of C, and represent each $c^{(k)}$ by $(c_1^{(k)}, c_2^{(k)}, ...)$. Let each X_n be the set of positive contractions in A, and define functions $f_n^{(k)} : X_n \to [0, \infty)$ by

$$\begin{aligned} f_n^{(1)}(x) &= \|(1 - x)d_n\|, \\ f_n^{(2)}(x) &= \|x\|_2, \\ f_n^{(2k+1)} &= \|\alpha^{g_k}(x) - x\| \qquad (k \in \mathbb{N}), \\ f_n^{(2k+2)} &= \|c_n^{(k)}x - xc_n^{(k)}\| \qquad (k \in \mathbb{N}). \end{aligned}$$

Given $m \in \mathbb{N}$ and $\varepsilon > 0$, there exists $\ell \in \mathbb{N}$ sufficiently large (along the filter ω) so that

$$e^{(\ell)} = (e_1^{(\ell)}, e_2^{(\ell)}, ...)$$

in A_ω satisfies

$$f_\omega^{(k)}(e_1^{(\ell)}, e_2^{(\ell)}, ...) < \varepsilon \qquad (1 \leq k \leq m).$$

Now the proof is finished by applying the ε-test. □

As in [16], we define a technical property called the *approximate Rokhlin property*, which would serve as a bridge connecting the W^*-Rokhlin property and the Rokhlin dimension.

Definition 4.3. Let A be a unital C^*-algebra and $\alpha : \mathbb{Z}^m \to \mathrm{Aut}(A)$ be a group action. We say α has the *approximate Rokhlin property* if for every $p \in \mathbb{N}$ there exist positive contractions $\{f_v\}_{v\in B_p}$ in $F(A)$ such that

(1) $(\alpha_\omega)^u(f_v) = f_{v+u} \mod p \qquad (v \in B_p,\ u \in \mathbb{Z}^m)$.
(2) $f_v f_u = 0 \qquad (v, u \in B_p,\ v \neq u)$.

(3) $1_A - \sum_{v\in B_p} f_v$ belongs to $J_A \cap F(A)$.
(4) $\sup_{k\in\mathbb{N}} \|1_A - f_0^k\|_{2,\omega} < 1$.

(here 0 is the origin in $\mathbb{Z}^m$).

With the aid of Lemma 4.2, it is not hard to pass from the W^*-Rokhlin property to the approximate Rokhlin property.

Lemma 4.4. *(cf.* [16, Propoition 5.8]*) Let A be a unital C^*-algebra and $\alpha : \mathbb{Z}^m \to \mathrm{Aut}(A)$ be a group action. Suppose the induced action $(\tilde{\alpha}, \bar{\alpha}) : \mathbb{Z}^m \to \mathrm{Aut}(\overline{A}^u, \partial_e T(A), E)$ has the W^*-Rokhlin property. Then α has approximate Rokhlin property.*

Proof. To simplify the notion, we write $(\mathcal{M}, K, E)$ for the triple $(\overline{A}^u, \partial_e T(A), E)$. Let $\{p_v\}_{v\in B_p}$ be projections in $\mathcal{M}^\omega \cap \mathcal{M}'$ satisfying the W^*-Rokhlin property. Recall that the canonical map $F(A) \to \mathcal{M}^\omega \cap \mathcal{M}'$ is surjective ([10]). Lifting the projections $\{p_v\}$ along this surjection, we obtain positive contractions $\{f'_v\}$ in $F(A)$ such that

(1) $(\alpha_\omega)^u(f'_v) - f'_{v+u} \in J_A \cap F(A) \mod p \qquad (v \in B_p,\ u \in \mathbb{Z}^m)$,
(2) $f'_v f'_u \in J_A \cap F(A) \qquad (u, v \in B_p,\ u \neq v)$,
(3) $1_A - \sum_v f'_v \in J_A \cap F(A)$.

Put $C := C^*(A, \{(\alpha_\omega)^u(f'_v) : u \in \mathbb{Z}^m,\ v \in B_p\})$. Then C is a separable C^*-subalgebra of A_ω satisfying $(\alpha_\omega)^u(C) = C$ for every $u \in \mathbb{Z}^m$. Since J_A is an α_ω-σ-ideal, there exists a positive contraction $u \in C' \cap J_A$ such that $(\alpha_\omega)^w(u) = u$ for all $w \in \mathbb{Z}^m$ and $uc = c$ for all $c \in C \cap J_A$. Define

$$f_v := (1-u)f'_v(1-u) \qquad (v \in B_p).$$

One checks that the family $\{f_v\}_{v\in B_p}$ has the desired properties. □

5. A Rokhlin type theorem

In this section we establish our Rokhlin type theorem for $\mathbb{Z}^m$-actions on C^*-algebras (Theorem 5.5). Let $\alpha : \mathbb{Z}^m \to \mathrm{Aut}(A)$ be a group action. The key observation, motivated by [20], is that suppose for every canonical generator α^i, we can find Rokhlin towers for α^i which are almost fixed by all the other generators $\alpha^1, ..., \alpha^{i-1}, \alpha^{i+1}, ..., \alpha^m$, then we can simply "multiply" the towers together and get Rokhlin towers for α. Here is the precise statement:

Proposition 5.1. *Let A be a unital C^*-algebra and $\alpha : \mathbb{Z}^m \to \mathrm{Aut}(A)$ be a group action. Suppose there exists $d \in \mathbb{N}$ such that for each $i \in \{1, ..., m\}$ the following holds: for every $\varepsilon > 0$, $p \in \mathbb{N}$, finite subset $F \subseteq A$, there exist positive contractions*

$$\{g_{i,k}^{(\ell)} : 0 \le k \le p-1,\ 0 \le \ell \le d\}$$

in A such that

(1) $\|\alpha^i(g_{i,k}^{(\ell)}) - g_{i,k+1}^{(\ell)}\| < \varepsilon \mod p \qquad (0 \le k \le p-1,\ 0 \le \ell \le d)$,
(2) $\|\alpha^j(g_{i,k}^{(\ell)}) - g_{i,k}^{(\ell)}\| < \varepsilon \qquad (1 \le j \le m,\ j \neq i,\ 0 \le k \le p-1,\ 0 \le \ell \le d)$,
(3) $\|[g_{i,k}^{(\ell)}, x]\| < \varepsilon \qquad (0 \le k \le p-1,\ 0 \le \ell \le d,\ x \in F)$,

(4) $\|g_{i,k}^{(\ell)} g_{i,n}^{(\ell)}\| < \varepsilon \qquad (0 \leq k \neq n \leq p-1,\ 0 \leq \ell \leq d)$,
(5) $\left\| \sum_{k,\ell} g_{i,k}^{(\ell)} - 1_A \right\| < \varepsilon$.

Then $\dim_{\mathrm{Rok}}(\alpha) \leq (d+1)^m - 1$.

Proof. Fix $\varepsilon > 0$, a finite subset $F \subseteq A$, and $p \in \mathbb{N}$. Let

$$\{g_{1,k}^{(\ell)} : 0 \leq k \leq p-1,\ 0 \leq \ell \leq d\}$$

be positive contractions in A satisfying (1) to (5) with respect to $\varepsilon' > 0$ and $F_1 := F$, where ε' is a small number to be determined later.

Define, for $i = 1, 2, ..., m-1$, inductively $F_{i+1} := F_i \cup \{g_{i,k}^{(\ell)}\}$ and let $\{g_{i+1,k}^{(\ell)} : 0 \leq k \leq p-1,\ 0 \leq \ell \leq d\}$ be positive contractions satisfying conditions (1) to (5) with respect to ε' and F_i. For $v = (v_1, ..., v_m) \in B_p$ and $(\ell_1, ..., \ell_m) \in \{0, ..., d\}^m$, define (positive contractions)

$$f_v^{(\ell_1,\ldots,\ell_m)} := \left(g_{m,v_m}^{(\ell_m)}\right)^{\frac{1}{2}} \cdots \left(g_{2,v_2}^{(\ell_2)}\right)^{\frac{1}{2}} \left(g_{1,v_1}^{(\ell_1)}\right) \left(g_{2,v_2}^{(\ell_2)}\right)^{\frac{1}{2}} \cdots \left(g_{m,v_m}^{(\ell_m)}\right)^{\frac{1}{2}}.$$

One checks that these elements form $(d+1)^m$ Rokhlin towers for α with respect to ε and F, provided that ε' is sufficiently small. □

For the rest of the section we will focus on establishing the conditions described in Proposition 5.1. One technical notion we need is the so-called *Property (SI)*; it was first introduced by Sato in [29], and later reformulated by Kirchberg and Rørdam in [10].

Definition 5.2. ([10, Definition 2.6]) Let A be a unital separable C^*-algebra with nonempty trace space. We say A has *property (SI)* if for all positive contractions $e, f \in F(A)$ with $e \in J_A$ and $\sup_{m\in\mathbb{N}} \|1_A - f^m\|_{2,\omega} < 1$, there exists an element $s \in F(A)$ with $fs = s$ and $s^*s = e$.

Lemma 5.3. *Let A be a unital separable C^*-algebra with $T(A) \neq \emptyset$, and $\alpha : \mathbb{Z}^m \to \mathrm{Aut}(A)$ be a group action. Suppose α has the approximate Rokhlin property and A has property (SI). Then for every $i \in \{1, ..., m\}$ and $p \in \mathbb{N}$ there exist $g_0, ..., g_{p-1}, v$ in $F(A)$ such that*

(i) $g_0, ..., g_{p-1}$ *are positive contractions,*
(ii) $g_k g_n = 0 \qquad (0 \leq k \neq n \leq p-1)$,
(iii) $(\alpha_\omega)^i(g_k) = g_{k+1} \mod p \qquad (0 \leq k \leq p-1)$,
(iv) $g_0 v = v$,
(v) $\sum_{k=0}^{p-1} g_k + v^*v = 1_A$,
(vi) $(\alpha_\omega)^{pi}(v) = v$,
(vii) $(\alpha_\omega)^j(g_k) = g_k \qquad (1 \leq j \neq i \leq m)$,
(viii) $(\alpha_\omega)^j(v) = v \qquad (1 \leq j \neq i \leq m)$.

Proof. The main idea is almost contained in [19, Proposition 4.5]. However, since the assumptions are slightly different, we include full details for the reader's convenience. Fix $i \in \{1, ..., m\}$ and $p \in \mathbb{N}$. First of all, since we are working with ultraproducts, it suffices

to find elements satisfying (i) to (viii) up to an arbitrarily small $\varepsilon > 0$. Let ℓ be a positive integer such that $\frac{2}{\sqrt{\ell p}} < \varepsilon$. By the approximate Rokhlin property, there exist $\{f_v\}_{v \in B_{\ell p}}$ satisfying

(1) $(\alpha_\omega)^u(f_v) = f_{v+u} \mod (\ell p) \qquad (v \in B_{\ell p},\ u \in \mathbb{Z}^m)$,
(2) $f_v f_u = 0 \qquad (v, u \in B_{\ell p},\ v \neq u)$,
(3) $e := 1_A - \sum_{v \in B_{\ell p}} f_v \in J_A \cap F(A)$,
(4) $\sup_n \|1_A - f_0^n\|_{2,\omega} < 1$.

Since A has property (SI), there is w in $F(A)$ such that

$$w^*w = e, \qquad \text{and} \qquad f_0 w = w.$$

In what follows, the coordinates of a point $v \in \mathbb{Z}^m$ will be denoted $v_1, v_2, ..., v_m$. Define, for $0 \leq k \leq \ell p - 1$,

$$g'_k := \sum_{\{v \in B_{\ell p} : v_i = k\}} f_v$$

and

$$s := \frac{1}{\sqrt{(\ell p)^{m-1}}} \sum_{\{u \in B_{\ell p} : u_i = 0\}} (\alpha_\omega)^u(w).$$

Then clearly we have

- $(\alpha_\omega)^j(g'_k) = g'_k \qquad (0 \leq k \leq \ell p - 1,\ 1 \leq j \neq i \leq m)$;
- $(\alpha_\omega)^i(g'_k) = g'_{k+1};\ \mod (\ell p) \qquad (0 \leq k \leq \ell p - 1)$;
- $\sum_{k=0}^{\ell p - 1} g'_k + e = 1_A$.

We make several more computations. First of all,

$$\begin{aligned}
s^*s &= \frac{1}{(\ell p)^{m-1}} \sum_{\{u \in B_{\ell p} : u_i = 0\}} \sum_{\{u' \in B_{\ell p} : u'_i = 0\}} \alpha_\omega^u(w^*) \alpha_\omega^{u'}(w) \\
&= \frac{1}{(\ell p)^{m-1}} \sum_{\{u \in B_{\ell p} : u_i = 0\}} \sum_{\{u' \in B_{\ell p} : u'_i = 0\}} \alpha_\omega^u(w^* f_0) \alpha_\omega^{u'}(f_0 w) \\
&= \frac{1}{(\ell p)^{m-1}} \sum_{\{u \in B_{\ell p} : u_i = 0\}} \sum_{\{u' \in B_{\ell p} : u'_i = 0\}} \alpha_\omega^u(w^*) f_u f_{u'} \alpha_\omega^{u'}(w) \\
&= \frac{1}{(\ell p)^{m-1}} \sum_{\{u \in B_{\ell p} : u_i = 0\}} \alpha_\omega^u(w^*) f_u f_u \alpha_\omega^u(w) \\
&= \frac{1}{(\ell p)^{m-1}} \sum_{\{u \in B_{\ell p} : u_i = 0\}} \alpha_\omega^u(w^*w) \\
&= e.
\end{aligned}$$

Second,

$$\begin{aligned} g_0' s &= \frac{1}{\sqrt{(\ell p)^{m-1}}} \sum_{\{v \in B_{\ell p} : v_i = 0\}} \sum_{\{u \in B_{\ell p} : u_i = 0\}} f_v (\alpha_\omega)^u(w) \\ &= \frac{1}{\sqrt{(\ell p)^{m-1}}} \sum_{\{v \in B_{\ell p} : v_i = 0\}} \sum_{\{u \in B_{\ell p} : u_i = 0\}} f_v f_u (\alpha_\omega)^u(w) \\ &= \frac{1}{\sqrt{(\ell p)^{m-1}}} \sum_{\{u \in B_{\ell p} : u_i = 0\}} f_u f_u (\alpha_\omega)^u(w) \\ &= \frac{1}{\sqrt{(\ell p)^{m-1}}} \sum_{\{u \in B_{\ell p} : u_i = 0\}} (\alpha_\omega)^u(w) \\ &= s. \end{aligned}$$

Finally, when $j \neq i$ we have,

$$\begin{aligned} \|(\alpha_\omega)^j(s) - s\| &= \frac{1}{\sqrt{(\ell p)^{m-1}}} \left\| \sum_{\{u \in B_{\ell p} : u_i = 0\}} (\alpha_\omega)^j (\alpha_\omega)^u(w) - \sum_{\{u \in B_{\ell p} : u_i = 0\}} (\alpha_\omega)^u(w) \right\| \\ &= \frac{1}{\sqrt{(\ell p)^{m-1}}} \left\| \sum_{\{u \in B_{\ell p} : u_i = 0, u_j = \ell p\}} (\alpha_\omega)^u(w) - \sum_{\{u \in B_{\ell p} : u_i = 0, u_j = 0\}} (\alpha_\omega)^u(w) \right\| \\ &\leq \frac{1}{\sqrt{(\ell p)^{m-1}}} \left(\left\| \sum_{\{u \in B_{\ell p} : u_i = 0, u_j = \ell p\}} (\alpha_\omega)^u(w) \right\| + \left\| \sum_{\{u \in B_{\ell p} : u_i = 0, u_j = 0\}} (\alpha_\omega)^u(w) \right\| \right). \end{aligned}$$

Since the terms in the norms are orthogonal, the C^*-identity implies that

$$\begin{aligned} \left\| \sum_{\{u \in B_{\ell p} : u_i = 0, u_j = \ell p\}} (\alpha_\omega)^u(w) \right\| &\leq \left\| \sum_{\{u \in B_{\ell p} : u_i = 0, u_j = \ell p\}} (\alpha_\omega)^u(w^* w) \right\|^{1/2} \leq \sqrt{(\ell p)^{m-2}} \cdot \|w^* w\|^{1/2} \\ &\leq \sqrt{(\ell p)^{m-2}} \end{aligned}$$

and a similar bound holds for the second norm. Therefore,

$$\|(\alpha_\omega)^j(s) - s\| \leq 2 \cdot \sqrt{\frac{(\ell p)^{m-2}}{(\ell p)^{m-1}}} = \frac{2}{\sqrt{\ell p}} < \varepsilon.$$

Since we work in the ultrapower, we may actually assume that

$$(\alpha_\omega)^j(s) = s \qquad (j \neq i).$$

Now define, for $0 \leq k \leq m-1$,

$$g_k := \sum_{r=0}^{\ell-1} g'_{k+rp}$$

and

$$v := \frac{1}{\sqrt{\ell}} \sum_{r=0}^{\ell-1} (\alpha_\omega)^{(rp)i}(s).$$

One checks that $g_0, g_1, ..., g_{m-1}, v$ satisfy the desired properties. □

Lemma 5.4. *Let A be a unital separable C^*-algebra with nonempty trace space $T(A)$, and $\alpha : \mathbb{Z}^m \to \mathrm{Aut}(A)$ be a group action. Suppose α has the approximate Rokhlin property and A has property (SI). Then for each $i \in \{1, ..., m\}$ and any $p \in \mathbb{N}$ there exist positive contractions*

$$a_0, ..., a_{p-1}, b_0, ..., b_{p-1}, c_0, ..., c_p$$

in $F(A)$ such that

(i) $(\alpha_\omega)^i(a_k) = a_{k+1}$ *and* $(\alpha_\omega)^i(b_k) = b_{k+1} \mod p \quad (0 \leq k \leq p-1)$.
(ii) $(\alpha_\omega)^i(c_k) = c_{k+1} \mod (p+1) \quad (0 \leq k \leq p)$.
(iii) $(\alpha_\omega)^j(a_k) = a_k$ *and* $(\alpha_\omega)^j(b_k) = b_k \quad (1 \leq j \neq i \leq m,\ 0 \leq k \leq p-1)$
(iv) $(\alpha_\omega)^j(c_k) = c_k \quad (1 \leq j \neq i \leq m,\ 0 \leq k \leq p)$
(v) $a_k a_n = b_k b_n = 0 \quad (0 \leq k \neq n \leq p-1)$.
(vi) $c_k c_n = 0 \quad (0 \leq k \neq n \leq p)$.
(vii) $b_k c_n = 0 \quad (0 \leq k \leq p-1,\ 0 \leq n \leq p)$.
(viii) $\sum_k a_k + \sum_k b_k + \sum_k c_k = 1_A$.

Proof. Copy the proof of [16, Theorem 6.4] verbatim, replacing [16, Proposition 6.2] by the previous lemma. Note that since all "matrix units" x_i's are fixed by $(\alpha_\omega)^j$, the entire image of $\psi : M_{\ell p+1}(\mathbb{C}) \to F(A)$ is fixed by $(\alpha_\omega)^j$ (here $j \neq i$). Therefore (iii) and (iv) are satisfied. □

Theorem 5.5. *Let A be a unital simple separable nuclear C^*-algebra with nonempty trace space $T(A)$, and let $\alpha : \mathbb{Z}^m \to \mathrm{Aut}(A)$ be a group action. Suppose*

(1) *A has property (SI),*
(2) *α is strongly outer,*
(3) *$T(A)$ is a Bauer simplex with finite dimensional extreme boundary,*
(4) *$\tau \circ \alpha = \tau$ for every $\tau \in T(A)$.*

Then $\dim_{\mathrm{Rok}}(\alpha) \leq 4^m - 1$.

Proof. Write $(\mathcal{M}, K, E)$ for the bundle $(\overline{A}^u, \partial_e T(A), E)$. Under the current assumptions, the induced group action $(\tilde{\alpha}, \bar{\alpha}) : \mathbb{Z}^m \to \mathrm{Aut}(\mathcal{M}, K, E)$ satisfies conditions (1) to (4) in Theorem 3.5. Therefore the action $(\tilde{\alpha}, \bar{\alpha})$ has the W^*-Rokhlin property. By Lemma 4.4, α has the approximate Rokhlin property. The previous lemma, together with [6, Proposition 2.8] (which converts towers of different heights to towers of the same height), shows that the assumptions of Proposition 5.1 are satisfied with $d = 3$. By the same proposition we have $\dim_{\mathrm{Rok}}(\alpha) \leq 4^m - 1$. □

Corollary 5.6. *Under the same assumptions as Theorem 5.5, the crossed product $A \rtimes_\alpha \mathbb{Z}^m$ is a unital simple separable C^*-algebra of finite nuclear dimension.*

Proof. From Theorem 5.5 and [30, Proposition 2.4], we see that the crossed product $A \rtimes_\alpha \mathbb{Z}^m$ has finite nuclear dimension. Simplicity follows from [12, Theorem 3.1]. □

6. Example: Bernoulli Actions

In this final section we study the Bernoulli actions. We will show that for a large class of C^*-algebras, the $\mathbb{Z}^m$-Bernoulli shift has finite Rokhlin dimension. Throughout the section all tensor products are understood as minimal tensor products.

We recall the definition of Bernoulli actions. Let A be a unital C^*-algebra and Γ be a discrete group. Define $\bigotimes_\Gamma A$ to be the inductive limit of the system $\{\bigotimes_F A : F \subseteq \Gamma \text{ finite subset}\}$, where the connecting maps are the natural inclusions. The (left) translation action of Γ on itself gives rise to an action $\sigma : \Gamma \to \mathrm{Aut}(\bigotimes_\Gamma A)$, called the *Bernoulli action* (or *Bernoulli shift*).

The following proposition is a slight variation of [29, Proposition 4.4].

Proposition 6.1. *Let A be a unital simple C^*-algebra with a unique trace. Suppose A is not isomorphic to the algebra of complex numbers $\mathbb{C}$. Then the Bernoulli action $\sigma : \mathbb{Z}^m \to \mathrm{Aut}(\bigotimes_{\mathbb{Z}^m} A)$ is strongly outer.*

Proof. For convenience we write $C := \bigotimes_{n\in\mathbb{Z}} A$. Let τ be the unique trace on C and let π_τ be the GNS representation of C associated to τ. Let $\tilde{\sigma} \in \mathrm{Aut}(\pi_\tau(C)'')$ be the weak extension of the action σ. We show that for every $v \in \mathbb{Z}^m \setminus \{0\}$, the automorphism $\tilde{\sigma}^v$ is not inner in $\pi_\tau(C)''$.

Assume on the contrary that $\tilde{\sigma}^v = \mathrm{Ad}\, W_0$ for some unitary W_0 in $\pi_\tau(C)''$. Note that $\tilde{\sigma}^{nv}(W_0) = W_0$ for all $n \in \mathbb{N}$. Fix $\varepsilon > 0$ and let x be a contraction in some finite tensor product $\bigotimes_F A \subseteq C$. There exist a finite subset $F' \subseteq \mathbb{Z}^m$ and an element $w_0 \in \bigotimes_{F'} A \subseteq C$ such that

$$\|W_0 - \pi_\tau(w_0)\|_{2,\tau} < \varepsilon.$$

Then

$$\|W_0 - \pi_\tau(\sigma^{nv}(w_0))\|_{2,\tau} < \varepsilon$$

for any $n \in \mathbb{N}$. If n is sufficiently large, then $\sigma^{nv}(w_0)$ commutes with x. Therefore W_0 commutes with $\pi_\tau(x)$ up to 2ε in the 2-norm. Since ε is arbitrary and τ is faithful (simplicity passes to inductive limits), we see that W_0 commutes with $\pi_\tau(x)$ exactly, and hence belongs to the center of $\pi_\tau(C)''$. Since C has a unique trace, $\pi_\tau(C)''$ is a factor (the center is trivial). This shows that $\mathrm{Ad}\, W_0$ is the identity map, a contradiction. □

Corollary 6.2. *Let A be either the matrix algebra $M_n(\mathbb{C})$ ($n \geq 2$) or the Jiang-Su algebra $\mathcal{Z}$ (see [7]), and let $\sigma : \mathbb{Z}^m \to \mathrm{Aut}(\bigotimes_{\mathbb{Z}^m} A)$ be the Bernoulli action. Then σ has finite Rokhlin dimension.*

Proof. In either case, A is a unital simple separable nuclear C^*-algebra with a unique trace. Therefore the previous proposition shows that the Bernoulli action σ is strongly outer. It is well-known that the infinite tensor product of a nuclear C^*-algebra remains to be nuclear. To apply Theorem 5.5, it remains to check that $\bigotimes_{\mathbb{Z}^m} A$ has property (SI). In the case that $A = M_n(\mathbb{C})$, the infinite tensor product is the UHF algebra of type n^∞, which is known to absorb the Jiang-Su algebra $\mathcal{Z}$ (for example, see [11] and [33]). In the case $A = \mathcal{Z}$, the infinite tensor product is isomorphic to $\mathcal{Z}$ itself by [7, Theorem 4], and in particular is Jiang-Su stable as well. Since Jiang-Su stability implies property (SI) [18, Theorem 1.1], the proof is complete. □

The following observation allows us to expand the class of C^*-algebra drastically.

Proposition 6.3. *Let A be a unital C^*-algebra and $B \subseteq A$ be a unital C^*-subalgebra (with $1_A \in B$). Let $\sigma : \mathbb{Z}^m \to \mathrm{Aut}(\bigotimes_{\mathbb{Z}^m} A)$ be the Bernoulli action. Then $\dim_{\mathrm{Rok}}(\sigma) \leq \dim_{\mathrm{Rok}}(\sigma|_{\bigotimes_{\mathbb{Z}^m} B})$.*

Proof. Assume $\dim_{\mathrm{Rok}}(\sigma|_{\bigotimes_{\mathbb{Z}^m} B}) = d < \infty$ otherwise there is nothing to prove. Fix a finite subset $\{a_1, ..., a_m\} \subseteq \bigotimes_{\mathbb{Z}^m} A$, $\varepsilon > 0$, and $n \in \mathbb{N}$. By approximation we may assume that $\{a_1, ..., a_m\} \subseteq \bigotimes_F A$ for some finite subset $F \subseteq \mathbb{Z}^m$. By assumption we can find positive contractions

$$\{f_v^{(\ell)}\}_{v \in B_n}^{\ell = 0,1,...,d}$$

in $\bigotimes_{\mathbb{Z}^m} B$ such that

(1) $\|\sigma^w(f_v^{(\ell)}) - f_{v+w}^{(\ell)}\| < \varepsilon' \mod n \qquad (0 \leq \ell \leq d,\ v \in B_n,\ w \in \mathbb{Z}^m)$;
(2) $\|f_v^{(\ell)} f_{v'}^{(\ell)}\| < \varepsilon' \qquad (0 \leq \ell \leq d,\ v, v' \in B_n,\ v \neq v')$.
(3) $\left\| \sum_{\ell=0}^d \sum_{v \in B_n} f_v^{(\ell)} - 1_A \right\| < \varepsilon'$.

for some small $\varepsilon' > 0$ to be determined later. Find a finite subset $F' \subseteq \mathbb{Z}^m$ and elements $\{g_v^{(\ell)}\}_{v \in B_n}^{\ell = 0,1,...,d}$ in $\bigotimes_{F'} B$ such that for $\ell = 0, 1, ..., d$ and $v \in B_n$,

$$\|g_v^{(\ell)} - f_v^{(\ell)}\| < \varepsilon'.$$

Then

$$\|\sigma^w(g_v^{(\ell)}) - g_{v+w}^{(\ell)}\| < 3\varepsilon' \mod n,$$

$$\left\| \sum_{v,\ell} g_v^{(\ell)} - 1_A \right\| \leq \left\| \sum_{v,\ell} (g_v^{(\ell)} - f_v^{(\ell)}) \right\| + \left\| \sum_{v,\ell} f_v^{(\ell)} - 1_A \right\| < [\mathrm{card}(B_n)(d+1) + 1]\varepsilon'$$

and

$$\begin{aligned} \|g_v^{(\ell)} g_{v'}^{(\ell)}\| &\leq \|[g_v^{(\ell)} - f_v^{(\ell)}] g_{v'}^{(\ell)}\| + \|f_v^{(\ell)} [g_{v'}^{(\ell)} - f_{v'}^{(\ell)}]\| + \|f_v^{(\ell)} f_{v'}^{(\ell)}\| \\ &< 3\varepsilon'. \end{aligned}$$

Observe that we can find $u \in \mathbb{Z}^m$ such that

$$[\sigma^u(g_v^{(\ell)}), a_j] = 0 \qquad (v \in B_n,\ 0 \leq \ell \leq d,\ 1 \leq j \leq m).$$

Define

$$h_v^{(\ell)} := \sigma^u(g_v^{(\ell)}) \qquad (v \in B_n,\ 0 \leq \ell \leq d).$$

Then

$$\|\sigma^w(h_v^{(\ell)}) - h_{v+w}^{(\ell)}\| = \|\sigma^{w+u}(g_v^{(\ell)}) - \sigma^u(g_{v+w}^{(\ell)})\| < \varepsilon'$$

and

$$\begin{aligned}\left\|\sum_{v,\ell} h_v^{(\ell)} - 1_A\right\| = \left\|\sum_{v,\ell} \sigma^u(g_v^{(\ell)}) - 1_A\right\| = \left\|\sum_{v,\ell} g_v^{(\ell)} - 1_A\right\| \\ < [\mathrm{card}(B_n)(d+1)+1]\varepsilon'.\end{aligned}$$

Finally,

$$\|h_v^{(\ell)} h_{v'}^{(\ell)}\| = \|g_v^{(\ell)} g_{v'}^{(\ell)}\| < 3\varepsilon'.$$

If ε' is chosen to be sufficiently small, then we have obtained the Rokhlin towers $\{h_v^{(\ell)}\}$ for σ. □

Example 6.4. Let A be a unital C^*-algebra. Assume one of the following holds:

(1) A contains a unital copy of the Jiang-Su algebra $\mathcal{Z}$; for example, when A is Jiang-Su stable.
(2) A contains a unital copy of a matrix algebra $M_n(\mathbb{C})$ for some $n \geq 2$; for exmaple, the set of all continuous matrix-valued functions $C(X, M_n(\mathbb{C}))$ on some compact Hausdorff space X.

Then the previous proposition, together with Corollary 6.2, implies that the Bernoulli action $\sigma : \mathbb{Z}^m \to \mathrm{Aut}(\bigotimes_{\mathbb{Z}^m} A)$ has finite Rokhlin dimension.

References

[1] Joan Bosa, Nathanial P. Brown, Yasuhiko Sato, Aaron Tikuisis, Stuart White, and Wilhelm Winter. Covering dimension of C^*-algebras and 2-coloured classification. *preprint, arxiv:1506.03974*.

[2] Alain Connes. Outer conjugacy classes of automorphisms of factors. *Ann. Sci. École Norm. Sup. (4)*, 8(3):383–419, 1975.

[3] Alain Connes. Classification of injective factors. Cases II_1, II_∞, III_λ, $\lambda \neq 1$. *Ann. of Math. (2)*, 104(1):73–115, 1976.

[4] George A. Elliott, Guihua Gong, Huaxin Lin, and Zhuang Niu. On the classification of simple amenable C^*-algebras with finite decomposition rank, II. *preprint, arxiv:1507.03437*.

[5] Richard H. Herman and Adrian Ocneanu. Stability for integer actions on UHF C^*-algebras. *J. Funct. Anal.*, 59(1):132–144, 1984.

[6] Ilan Hirshberg, Wilhelm Winter, and Joachim Zacharias. Rokhlin dimension and C^*-dynamics. *Comm. Math. Phys.*, 335(2):637–670, 2015.

[7] Xinhui Jiang and Hongbing Su. On a simple unital projectionless C^*-algebra. *Amer. J. Math.*, 121(2):359–413, 1999.

[8] Shizuo Kakutani. Induced measure preserving transformations. *Proc. Imp. Acad. Tokyo*, 19:635–641, 1943.

[9] Eberhard Kirchberg. The classification of purely inifinite C^*-algebras using Kasparov's theory. *to appear in the Fields Institute Communication series.*

[10] Eberhard Kirchberg and Mikael Rørdam. Central sequence C^*-algebras and tensorial absorption of the Jiang-Su algebra. *J. Reine Angew. Math.*, 695:175–214, 2014.

[11] Eberhard Kirchberg and Wilhelm Winter. Covering dimension and quasidiagonality. *Internat. J. Math.*, 15(1):63–85, 2004.

[12] Akitaka Kishimoto. Outer automorphisms and reduced crossed products of simple C^*-algebras. *Comm. Math. Phys.*, 81(3):429–435, 1981.

[13] Akitaka Kishimoto. The Rohlin property for automorphisms of UHF algebras. *J. Reine Angew. Math.*, 465:183–196, 1995.

[14] Akitaka Kishimoto. The Rohlin property for shifts on UHF algebras and automorphisms of Cuntz algebras. *J. Funct. Anal.*, 140(1):100–123, 1996.

[15] Akitaka Kishimoto. Automorphisms of A$\mathbf{T}$ algebras with the Rohlin property. *J. Operator Theory*, 40(2):277–294, 1998.

[16] Hung-Chang Liao. A Rokhlin type theorem for simple C^*-algebras of finite nuclear dimension. *J. Funct. Anal.*, 270(10):3675–3708, 2016.

[17] Hiroki Matui and Yasuhiko Sato. $\mathcal{Z}$-stability of crossed products by strongly outer actions. *Comm. Math. Phys.*, 314(1):193–228, 2012.

[18] Hiroki Matui and Yasuhiko Sato. Strict comparison and $\mathcal{Z}$-absorption of nuclear C^*-algebras. *Acta Math.*, 209(1):179–196, 2012.

[19] Hiroki Matui and Yasuhiko Sato. $\mathcal{Z}$-stability of crossed products by strongly outer actions II. *Amer. J. Math.*, 136(6):1441–1496, 2014.

[20] Hideki Nakamura. The Rohlin property for $\mathbb{Z}^2$-actions on UHF algebras. *J. Math. Soc. Japan*, 51(3):583–612, 1999.

[21] Adrian Ocneanu. A Rohlin type theorem for groups acting on von Neumann algebras. In *Topics in modern operator theory (Timişoara/Herculane, 1980)*, volume 2 of *Operator Theory: Adv. Appl.*, pages 247–258. Birkhäuser, Basel-Boston, Mass., 1981.

[22] Adrian Ocneanu. *Actions of discrete amenable groups on von Neumann algebras*, volume 1138 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, 1985.

[23] Hiroyuki Osaka and N. Christopher Phillips. Furstenberg transformations on irrational rotation algebras. *Ergodic Theory Dynam. Systems*, 26(5):1623–1651, 2006.

[24] Hiroyuki Osaka and N. Christopher Phillips. Stable and real rank for crossed products by automorphisms with the tracial Rokhlin property. *Ergodic Theory Dynam. Systems*, 26(5):1579–1621, 2006.

[25] Narutaka Ozawa. Dixmier approximation and symmetric amenability for C*-algebras. *J. Math. Sci. Univ. Tokyo*, 20(3):349–374, 2013.

[26] N. Christopher Phillips. A classification theorem for nuclear purely infinite simple C^*-algebras. *Doc. Math.*, 5:49–114 (electronic), 2000.

[27] Vladimir A. Rohlin. A "general" measure-preserving transformation is not mixing. *Doklady Akad. Nauk SSSR (N.S.)*, 60:349–351, 1948.

[28] Mikael Rørdam. The stable and the real rank of $\mathcal{Z}$-absorbing C^*-algebras. *Internat. J. Math.*, 15(10):1065–1084, 2004.

[29] Yasuhiko Sato. The Rohlin property for automorphisms of the Jiang-Su algebra. *J. Funct. Anal.*, 259(2):453–476, 2010.

[30] Gábor Szabó. The Rokhlin dimension of topological $\mathbb{Z}^m$-actions. *Proc. Lond. Math. Soc. (3)*, 110(3):673–694, 2015.

[31] Gábor Szabó, Jianchao Wu, and Joachim Zacharias. Rokhlin dimension for actions of residually finite groups. *preprint, arxiv:1408.6096.*

[32] Aaron Tikuisis, Stuart White, and Wilhelm Winter. Quasidiagonality of nuclear C^*-algebras. *preprint, arxiv:1509.08318*.
[33] Wilhelm Winter. Decomposition rank and $\mathcal{Z}$-stability. *Invent. Math.*, 179(2):229–301, 2010.
[34] Wilhelm Winter. Nuclear dimension and $\mathcal{Z}$-stability of pure C^*-algebras. *Invent. Math.*, 187(2):259–342, 2012.
[35] Wilhelm Winter and Joachim Zacharias. Completely positive maps of order zero. *Münster J. Math.*, 2:311–324, 2009.